



# Defects in the COG complex and COG-related trafficking regulators affect neuronal Golgi function

Leslie K. Climer<sup>1</sup>, Maxim Dobretsov<sup>2</sup> and Vladimir Lupashin<sup>1\*</sup>

<sup>1</sup> Department of Physiology and Biophysics, College of Medicine, University of Arkansas for Medical Sciences, Little Rock, AR, USA, <sup>2</sup> Department of Anesthesiology, College of Medicine, University of Arkansas for Medical Sciences, Little Rock, AR, USA

## OPEN ACCESS

### Edited by:

Catherine Rabouille,  
Hubrecht Institute, Netherlands

### Reviewed by:

Wim Annaert,  
Catholic University of Leuven, Belgium  
Rainer Duden,  
University of Luebeck, Germany

### \*Correspondence:

Vladimir Lupashin  
vlupashin@uams.edu

### Specialty section:

This article was submitted to  
Neurodegeneration,  
a section of the journal  
Frontiers in Neuroscience

**Received:** 14 August 2015

**Accepted:** 12 October 2015

**Published:** 27 October 2015

### Citation:

Climer LK, Dobretsov M and  
Lupashin V (2015) Defects in the COG  
complex and COG-related trafficking  
regulators affect neuronal Golgi  
function. *Front. Neurosci.* 9:405.  
doi: 10.3389/fnins.2015.00405

The Conserved Oligomeric Golgi (COG) complex is an evolutionarily conserved hetero-octameric protein complex that has been proposed to organize vesicle tethering at the Golgi apparatus. Defects in seven of the eight COG subunits are linked to Congenital Disorders of Glycosylation (CDG)-type II, a family of rare diseases involving misregulation of protein glycosylation, alterations in Golgi structure, variations in retrograde trafficking through the Golgi and system-wide clinical pathologies. A troublesome aspect of these diseases are the neurological pathologies such as low IQ, microcephaly, and cerebellar atrophy. The essential function of the COG complex is dependent upon interactions with other components of trafficking machinery, such as Rab-GTPases and SNAREs. COG-interacting Rabs and SNAREs have been implicated in neurodegenerative diseases like Alzheimer's disease and Parkinson's disease. Defects in Golgi maintenance disrupts trafficking and processing of essential proteins, frequently associated with and contributing to compromised neuron function and human disease. Despite the recent advances in molecular neuroscience, the subcellular bases for most neurodegenerative diseases are poorly understood. This article gives an overview of the potential contributions of the COG complex and its Rab and SNARE partners in the pathogenesis of different neurodegenerative disorders.

**Keywords:** conserved oligomeric Golgi complex, COG, congenital disorders of glycosylation, neurodegeneration, glycosylation, vesicular trafficking, Rab, SNARE

## INTRODUCTION

The Conserved Oligomeric Golgi (COG) complex is an evolutionarily conserved hetero-octameric protein complex that is a proposed membrane tether during vesicular trafficking at the Golgi apparatus (Lupashin and Ungar, 2008; Reynders et al., 2011; Miller and Ungar, 2012; Willett et al., 2013b). COG is composed of two functionally distinct sub-complexes lobe A (COG1-4) and lobe B (COG5-8) (Fotso et al., 2005; Ungar et al., 2005). Secretory and transmembrane proteins make up 30–50% of all cellular proteins, and are trafficked through the endoplasmic reticulum (ER) to the Golgi for folding and modifications before delivery to their final destination. Secretory cargo molecules are thought to travel through the Golgi complex mostly inside flat cisternae that are constantly maturing in a *cis-to-trans* (anterograde) fashion via the so called cisternal maturation mechanism (Glick and Malhotra, 1998). However, resident Golgi proteins and Soluble NSF Attachment protein Receptors (SNAREs) are constantly recycled back in vesicular

carriers to replenish the content of newly formed cisternae. COG regulates the recycling of vesicles containing glycosylation enzymes and other resident Golgi proteins in a *trans*-to-*cis* (retrograde) direction. An intricate assortment of trafficking machineries including small Rab-GTPases, SNAREs, Sec1/Munc18 (SM) proteins, vesicular coat proteins, and tethering proteins are required for vesicular transport (Bonifacino and Glick, 2004). Intracellular pathways rely on these protein families at each step of vesicular transport. Though the functional interaction between the tethers and other trafficking regulators is not completely understood, a multi-subunit tethering complex (MTC), like the COG complex, may coordinate the interactions between all other components of the trafficking machinery at the site of vesicle docking on the target membrane for efficient fusion of the two membranes (Cottam and Ungar, 2012; Willett et al., 2013b).

SNAREs are an essential COG partner. SNAREs catalyze the fusion of the vesicle membrane with the target membrane by the assembly of a quaternary SNARE complex that functions as a zipper to coalesce the opposing lipid bilayers. While SNAREs alone have an innate ability to fuse membranes, the SNARE regulatory proteins and tethering complexes are thought to be necessary for physiologically relevant fusion events (Rizo and Südhof, 2012). The COG complex interacts with at least two Golgi SNARE complexes: the *cis*-Golgi STX5/GOSR1(GS28)/YKT6/BET1L(GS15) complex; and the *trans*-Golgi STX16/STX6/VTI1a/VAMP4 complex (Shestakova et al., 2007; Laufman et al., 2009, 2013b). Additionally, SNARE complexes that contain Sec22b, GOSR2(GS27), or SNAP29 are evidenced to interact with COG (Kudlyk et al., 2013; Willett et al., 2013a). SNARE partner transitions connect all compartments of the endocytic and secretory pathways. Thus, it is likely that a defect in one trafficking step may have a cascading effect on all other cargo trafficking steps. In this review we will focus on those SNAREs where deficiency was shown to be associated with neuronal abnormalities, i.e., Ykt6, Sec22b, STX5, SNAP29, GS27, GS28, and Vti1a/b (Table 1).

The essential function of the COG complex has been proposed to depend not only upon SNAREs interactions, but also interactions with Rabs (Miller et al., 2013; Willett et al., 2013a; Figure 1). Within this context, Rabs are believed to act as molecular switches that cycle between GDP-bound (inactive) and GTP-bound (active) states and regulate cargo trafficking by acting in an intracellular compartment-specific manner. Active (GTP-bound) Rabs regulate trafficking by binding effector molecules like tethering factors and motor proteins (D'Adamo et al., 2014). Binding to some effectors leads to the activation of other Rabs, in a sequence known as the Rab Cascade (Pfeffer, 2013). Nine distinct, Golgi-localized Rabs are known to interact with COG (Miller et al., 2013; Willett et al., 2013b, 2014). The compartment-specific nature of these different Rabs makes them a potential landmark for COG membrane localization and interaction. As with the SNAREs, this review will limit the discussion to COG partners Rab1a, Rab1b, Rab2, Rab4a, and Rab6a that are implicated in neuronal abnormalities (Table 1).

## NEUROPHYSIOLOGICAL ABNORMALITIES AND COG DEFECTS

Since 2004, defects in seven out of the eight COG subunits have been associated with human Congenital Disorders of Glycosylation (CDG)-type II, a growing family of diseases involving malfunctions in the processing of N- and O-linked glycans and resulting from mutations in proteins involved with glycosylation (Wu et al., 2004; Spaapen et al., 2005; Foulquier et al., 2006, 2007; Kranz et al., 2007; Ng et al., 2007, 2011; Paesold-Burda et al., 2009; Reynders et al., 2009; Richardson et al., 2009; Lübbehuisen et al., 2010; Fung et al., 2012; Huybrechts et al., 2012; Rymen et al., 2012, 2015; Kodera et al., 2015; Table 1). Glycosylation is a highly dynamic process that occurs in the ER and Golgi which requires an estimated two percent of the human genome to encode the enzymes and trafficking components for the proper maturation of newly formed glycan chains (Freeze et al., 2014). COG deficiency can cause a redistribution of COG-dependent Golgi resident proteins, including glycosylation enzymes. Most COG-CDG patients have defects in sialylation and galactosylation, as indicated by fluorescent lectin staining of plasma membrane glycoconjugates from patient fibroblasts and MALDI-TOF mass spectrometry of serum glycoproteins (Foulquier et al., 2007; Kranz et al., 2007; Paesold-Burda et al., 2009; Reynders et al., 2009; Zeevaert et al., 2009a,b). Along with other multi-system pathologies, COG-CDG patients display mild to severe neurological defects including hypotonia, intellectual disability, developmental delays, epilepsy, and ataxia (Table 1). Specific symptoms and the severity of condition appears to relate to the COG subunit that is deficient with COG6 and COG7 patients demonstrating the most severe phenotypes (Rymen et al., 2015). Additionally, COG defects have not yet been attributed to any other subtype of CDG.

Several CDGs result from mutated COG subunits that are either severely truncated or rapidly degraded. Loss of one COG subunit can destabilize the remaining subunits and reduce their expression and association with the Golgi. Early studies of the COG3 subunit invoked participation of the COG complex in the proper distribution of Golgi enzymes. COG3 depletion by siRNA in HeLa cells causes extensive Golgi fragmentation and destabilization of the COG complex (Zolov and Lupashin, 2005). COG3 and COG7 knockdown generates an accumulation of COG complex-dependent (CCD) vesicles carrying the SNAREs GS15 and GS28, and Golgi enzymes MAN2A1 and GALNT2 (Zolov and Lupashin, 2005; Shestakova et al., 2006). The accumulation of CCD vesicles suggests that in COG deficient cells a significant fraction of Golgi glycosylation enzymes are separated from the proteins they need to modify. COG8-CDG patient fibroblasts have decreased levels of the other lobe B subunits (COG5, COG6, and COG7) all of which have lost their association with the Golgi (Foulquier et al., 2007; Kranz et al., 2007). COG lobe B destabilization was also seen in COG7-CDG patient fibroblasts, resulting in the loss of COG6 association with the Golgi (Kudlyk et al., 2013). The loss of COG also challenges the function of interacting SNAREs. The endosome-to-*trans*-Golgi Network (TGN) SNARE protein STX16 was mislocalized in COG8-CDG patient fibroblasts (Willett et al., 2013a), and the


**TABLE 1 | Neurological phenotypes associated with COG and COG-interacting Rabs and SNAREs.**

Protein	Disorder	Associated neurological manifestation	References
<b>COG PROTEINS</b>			
COG1	CDG-IIg (COG1-CDG)	Cerebral atrophy, developmental delay, hypotonia	Foulquier et al., 2006
COG2	CDG-II (COG2-CDG)	Developmental delay, epilepsy	Kodera et al., 2015
COG4	CDG-IIj (COG4-CDG)	Developmental delay, epilepsy, hypotonia, lack of speech, nystagmus	Reynders et al., 2009; Ng et al., 2011
COG5	CDG-IIi (COG5-CDG)	Ataxia, cerebral atrophy, developmental delay, epilepsy, hypotonia	Paesold-Burda et al., 2009; Fung et al., 2012; Rymen et al., 2012
COG6	CDG-III (COG6-CDG) Shaheen syndrome (SHNS)	Ataxia, cerebral atrophy, developmental delay, epilepsy, hypotonia, optic nerve atrophy, sensorineural hearing loss Intellectual disability	Lübbhusen et al., 2010; Huybrechts et al., 2012; Shaheen et al., 2013; Rymen et al., 2015
COG7	CDG-IIe (COG7-CDG)	Cerebral atrophy, developmental delay, epilepsy, hypotonia	Wu et al., 2004; Morava et al., 2007; Ng et al., 2007; Zeevaert et al., 2009a
COG8	CDG-IIh (COG8-CDG)	Cerebral atrophy, developmental delay, hypotonia	Foulquier et al., 2007; Kranz et al., 2007
<b>SNARE PROTEINS</b>			
Ykt6	Parkinson's Disease	Trafficking defects and cytotoxicity <i>in vitro</i> in NRK and PC12 cell lines	Hasegawa et al., 2003, 2004
Sec22b	Parkinson's Disease	Trafficking defects and cytotoxicity <i>in vitro</i> in NRK and PC12 cell lines	Hasegawa et al., 2003, 2004
STX5	Parkinson's Disease Alzheimer's Disease	Trafficking defects and cytotoxicity <i>in vitro</i> in NRK and PC12 cell lines Regulates processing of APP in PC12, HeLa, COS-7, and NG108-15 cell lines and primary cultures of rat hippocampal neurons	Suga et al., 2005b; Thayanidhi et al., 2010; Rendón et al., 2013; Suga et al., 2015
SNAP29	CEDNIK-Neuro-cutaneous syndrome	Cerebral Dysgenesis, Neuropathy, Ichthyosis, and Keratoderma	Sprecher et al., 2005; Fuchs-Telem et al., 2011
GS27	Myoclonus epilepsy/early ataxia Parkinson's Disease	Action myoclonus, mild cerebral atrophy, and early ataxia Trafficking defects and cytotoxicity <i>in vitro</i> in NRK and PC12 cell lines	Thayanidhi et al., 2010; Corbett et al., 2011
GS28	Neurodegeneration	Retinal degeneration in <i>in vivo Drosophila</i> photoreceptors	Rosenbaum et al., 2014
Vti1a/b	Neurodegeneration	Perinatal lethality in double knockouts in an <i>in vivo</i> mouse model. Neuronal axon tracks missing, reduced in size or misrouted	Kunwar et al., 2011; Walter et al., 2014
<b>RAB PROTEINS</b>			
Rab1a	Parkinson's Disease	Neuroprotective in <i>C. elegans</i> , <i>D. melanogaster</i> and primary rat neuron cultures. Rescue from the neurotoxic effects of $\alpha$ -synuclein	Cooper et al., 2006; Gitler et al., 2008
Rab1b	Alzheimer's Disease	Dominant negative mutant of Rab1b blocks trafficking of APP and decreases the secretion of A $\beta$	Dugan et al., 1995
Rab2	Parkinson's Disease	Reduced expression of Rab2 can rescue Golgi fragmentation in PD models	Rendón et al., 2013
Rab4a	Neumann-Pick disease Alzheimer's Disease Down's syndrome	Reduced Rab4-dependent recycling <i>in vitro</i> in Neumann-Pick type A and type C fibroblasts. Postmortem samples: increased Rab4 in patients with AD and mild cognitive disorder A $\beta$ partially co-localizes with Rab4 in a mouse model of Down Syndrome	Cataldo et al., 2000; Choudhury et al., 2004; Arriagada et al., 2010; Ginsberg et al., 2010
Rab6a	Alzheimer's Disease	Dominant negative mutant of Rab6 increases the secretion of soluble APP and decreased A $\beta$ secretion	McConlogue et al., 1996

STX5/GS28/Ykt6/GS15 and STX6/STX16/Vti1a/VAMP4 SNARE complexes were destabilized in both COG7- and COG8-CDG patient fibroblasts (Laufman et al., 2013a). In a non-CDG patient presenting intellectual disability, Shaheen et al. identified a mutation in COG6 which resulted in reduced COG6 and STX6 protein expression (Shaheen et al., 2013). Anterograde trafficking does not appear to be affected in cells with COG mutations, but retrograde trafficking is affected as indicated by a partial resistance to treatment with the transport inhibitor brefeldin A (Steet and Kornfeld, 2006; Foulquier et al., 2007; Kranz et al., 2007; Ng et al., 2007; Paesold-Burda et al., 2009; Reynders et al., 2009) and by endosome-to-TGN trafficking defects elucidated by application of Shiga toxin and SubAB toxin (Zolov and Lupashin, 2005; Smith et al., 2009). Therefore, retrograde intra-Golgi and endosome-to-TGN sorting are particularly impaired by COG deficiency.

## NEUROPATHOLOGY AND DEFECTS IN COG-ASSOCIATED PROTEINS

Extensive *in vitro* analyses in control and disease models demonstrate that genetic deficiency in SNARE and Rab COG partners may also result in disintegration of the Golgi apparatus, thereby potentially influencing neurological impairment (Table 1). Interestingly, the therapeutic implications of studying neurodegeneration associated with defective COG function is broader than the COG-CDG patient population (D'Adamo et al., 2014; Rymen et al., 2015). Golgi fragmentation is a common feature of neurodegenerative diseases (Gonatas et al., 2006). Current theories argue that the Golgi fragmentation seen in Alzheimer's Disease (AD) and Parkinson's Disease (PD) is either a result of misfolded or aggregated proteins, or that fragmented Golgi causes etiologically important proteins



**FIGURE 1 | Model of COG trafficking within neurons.** A cartoon depicting the proposed roles for the COG complex (depicted as red shapes for lobe A and green shapes for lobe B subcomplexes) and its interacting protein partners in Golgi trafficking: Rabs (yellow circles), and SNAREs (purple lines). Right panel: Immunofluorescence images of COG complex subunit COG3 (top-red) and Golgi SNARE GS15 (middle-green) in rat dorsal root ganglion. Perinuclear (DAPI) co-localization is indicated by yellow in the merged image (bottom). Scale Bar = 10  $\mu$ m.

to aggregate and misfold leading to further progression of these diseases (Gonatas et al., 2006; Nakagomi et al., 2008; Bellouze et al., 2014; Joshi and Wang, 2015). However, Golgi fragmentation has also been linked to SNARE and Rab proteins making it difficult to pinpoint a single disease progression. Thus, the COG-Rab-SNARE dynamic is important for understanding neurodegenerative phenotypes.

## SNARES

Genetic deficiency in SNAREs was shown to be associated with the progression of neurodegenerative diseases like AD and PD. PD is marked by the presence of Lewy bodies which are principally composed of aggregated  $\alpha$ -synuclein. Under physiological conditions,  $\alpha$ -synuclein may regulate vesicle trafficking and promote synaptic transmission by binding directly to SNAREs and stimulating SNARE complex formation (Burré et al., 2010; Thayanidhi et al., 2010). Overexpression of wild type or the PD-associated mutant of  $\alpha$ -synuclein (A53T) leads to cytotoxicity and inhibition of ER-to-Golgi trafficking in *in vitro* models which can be partially suppressed by

co-overexpression of SNAREs Ykt6 or Sec22. Ykt6—a protein enriched in neurons—was more protective than Sec22 and suggests a specialized role in mammals (Hasegawa et al., 2003, 2004). *In vitro* binding experiments also point toward the direct interaction of  $\alpha$ -synuclein with STX5 and GS27 along with mutant  $\alpha$ -synuclein which destabilizes the STX5-GS27-rbet1-sec22b SNARE complex (Thayanidhi et al., 2010). This line of evidence further implicates trafficking defects in bringing about neurodegenerative cytotoxicity.

Although the exact molecular mechanisms connecting Golgi fragmentation and disease mutations is still much under investigation, *in vitro* models can be used to recapitulate Golgi fragmentation seen in neurodegenerative disorders (Suga et al., 2005a). Fragmentation has been recreated *in vitro* and can be rescued by regulating levels of SNAREs. A STX5 knockdown is known to induce Golgi fragmentation in HeLa cells and cultured neurons (Suga et al., 2005a; Amessou et al., 2007). On the other hand, in PC12 cells treated with 6-hydroxydopamine or methamphetamine (an established *in vitro* PD model), a decrease in the level of STX5 rescues Golgi fragmentation (Rendón et al., 2013). This same study also demonstrated



that Golgi fragmentation could intensify disease progression by inducing  $\alpha$ -synuclein aggregation and the formation of Lewy bodies.

STX5 defects have also been shown to affect processing of AD-related proteins. In AD, amyloid precursor protein (APP) undergoes a series of proteolytic events by  $\beta$ - and  $\gamma$ -secretases to create the amyloidogenic variants of  $\beta$ -amyloid ( $A\beta$ ) that are longer and more likely to form aggregates (Peric and Annaert, 2015). Presenilins form a complex with  $\gamma$ -secretase, and mutations in presenilin 1 (PS1) are the most frequently associated mutations found in AD which result in increased production of  $A\beta$ , or altered ratios of amyloid peptide species (Hardy, 2006; Saito et al., 2011; De Strooper et al., 2012). Overexpressed STX5 was shown to co-localize with and directly bind to PS1. Further, STX5 overexpression increased the accumulation of APP in the ER and cis-Golgi and inhibited  $A\beta$  secretion in a neuroblastoma cell line (NG108-15) (Suga et al., 2005b). In NG108-15 cells expressing the familial AD mutation PS1 $\Delta$ E9, STX5 was shown to have a decreased association with presenilin. A study of ER stress in an AD model demonstrated that ER stressors can increase synthesis of STX5 and its accumulation in the ER-Golgi intermediate compartment (ERGIC) and transport vesicles. Thus, upregulation of trafficking machinery induced by ER stress could be a cellular mechanism for correcting the accumulation of the amyloidogenic cleavage products of APP (Suga et al., 2015).

As stated above, the cis-Golgi SNARE GS27 (GOSR2) binds to  $\alpha$ -synuclein and is part of a SNARE complex that is destabilized by mutant  $\alpha$ -synuclein. GS27 has also been shown to be associated with a neurological disorder in humans. Six patient were identified bearing a mutation that results in improper subcellular localization and loss of function of GS27 leading to symptoms common in COG-CDG patients such as cerebral atrophy, epilepsy, and early ataxia (Corbett et al., 2011; **Table 1**).

GS28 (GOSR1) is a Golgi SNARE involved in both ER-to-Golgi and intra-Golgi transport (Nagahama et al., 1996; Subramaniam et al., 1996), and accordingly has been shown to be associated with three SNARE complexes (Zhang and Hong, 2001; Parlati et al., 2002; Xu et al., 2002; Siddiqi et al., 2010). GS28 mutants have been used to study retinal degeneration in *Drosophila* photoreceptors (Rosenbaum et al., 2014). Lack of expression of GS28 in mutant flies alters trafficking and glycosylation of rhodopsin (Rh1). The photoreceptors in these mutants also exhibit enlarged ER and Golgi membranes and retinal degeneration over time.

Vt1a is a TGN-localized SNARE that functions in vesicle generation and Ca<sup>2+</sup> channel trafficking (von Mollard et al., 1997; Lupashin et al., 1997; Walter et al., 2014). A double knockout of Vt1a and Vt1b genes results in progressive neurodegeneration and perinatal lethality in a mouse model (Kunwar et al., 2011). Single knockout of Vt1a or Vt1b does not result in a lethal phenotype indicating overlapping functions of these proteins. The death-after-birth clearly demonstrates that these SNAREs are not required during organismal development, indicating a specialized requirement

for Vt1a and Vt1b in neurons leading to neurodegeneration in the double-knockout animals (Walter et al., 2014). Vt1a SNARE partners, STX6 and STX16, are also required for neurite outgrowth (Chua and Tang, 2008; Kabayama et al., 2008).

In humans, a disease known as CEDNIK (Cerebral Dysgenesis, Neuropathy, Ichthyosis, and Keratoderma) syndrome has been linked with loss of function mutations in SNAP29 (Sprecher et al., 2005; Fuchs-Telem et al., 2011). SNAP29 is a member of the SNAP25 family that localizes to the Golgi, endosomal, and lysosomal compartments (Steegmaier et al., 1998). CEDNIK patients exhibit severe neuropathy likely due to the loss of SNAP29 functional involvement in neurotransmission (Pan et al., 2005) and trafficking within neuroglia during active myelination (Schardt et al., 2009).

## RABS

As for SNAREs, regulating levels of Rabs has been shown to rescue Golgi fragmentation in multiple *in vitro* models of AD and PD. For example, overexpression of Rab1 can rescue Golgi fragmentation while reduced expression of Rab2 has the same affect in PD models which demonstrates the delicate balance in the regulatory functions of Golgi-associated Rabs (Rendón et al., 2013). Overexpression of Rab1 was shown to be neuroprotective in *Caenorhabditis elegans*, *Drosophila melanogaster* and primary rat neuron cultures (Cooper et al., 2006; Gitler et al., 2008). Rab1 is a key protein in maintaining Golgi architecture and function (Haas et al., 2007). It can also promote the restoration of ER-to-Golgi trafficking and thus afford rescue from the neurotoxic effects of  $\alpha$ -synuclein (Cooper et al., 2006; Gitler et al., 2008).

Multiple Rabs are associated with processing of APP. For example, ERGIC and cis-Golgi Rab1b-dependent trafficking could modulate the processing of APP as demonstrated in an *in vitro* system in which a dominant-negative mutant of Rab1b blocked trafficking of APP and decreased the secretion of  $A\beta$  (Dugan et al., 1995). trans-Golgi Rab6A is also implicated in APP trafficking. The dominant negative mutant of Rab6 increased the secretion of soluble APP and decreased  $A\beta$  secretion (McConlogue et al., 1996). Rab6 membrane association is dependent upon PS1 based on studies using PS1 knockout fibroblasts, thus implicating PS1 in vesicular trafficking (Scheper et al., 2004).

Golgi-associated and endosomal Rab4 is important for a specific pool of constitutively recycling endosomes which are apparently critical for dendritic spine size. Rab4-dependent recycling is greatly reduced in fibroblasts of patients with type A/B Niemann-Pick disease, a sphingolipid storage disorder which commonly manifests with neurological symptoms such as developmental delay and dementia (Choudhury et al., 2004). Postmortem samples revealed that Rab4 is upregulated in patients with AD and mild cognitive disorder (Cataldo et al., 2000; Ginsberg et al., 2010), and  $A\beta$  is known to partially co-localize within Rab4 positive compartments in a mouse model of Down Syndrome (Arriagada et al., 2010) indicating that defects in endosomal sorting may underpin these disorders (Peric and Annaert, 2015).

## CONCLUSIONS AND PERSPECTIVES

The COG complex and its Rab and SNARE partners are evolutionarily conserved and ubiquitously expressed across multiple tissues and organ systems in humans. Yet, neurological symptoms are the most debilitating and troublesome clinical manifestations of COG-associated disorders. Why is this the case? We propose that brain-specific manifestations of COG defects result from either COG-dependent (a) glycosylation defects and/or, (b) trafficking defects and/or, (c) a yet unknown neuron/neuroglia-specific function of the COG complex and its partners.

Neuronal function critically depends on coordinated delivery of properly modified ion channels, transporters and components of the synaptic apparatus at the appropriate rates and over long distances, to specific subcellular compartments. Remarkably, the localization of the synaptobrevin homolog Snc1 is altered in COG deficient yeast cells (Whyte and Munro, 2002). In addition, underglycosylated low density lipoprotein receptor is severely destabilized in CHO cells deficient for COG1 or COG2 proteins (Kingsley et al., 1986). Since glycosylation of channels, transporters and transport regulators is essential for their correct delivery, stability and/or activity (Gong et al., 2002; Watanabe et al., 2004; Scott and Panin, 2014), it is reasonable to predict that a majority of underglycosylated ion channels and transporters may similarly be destabilized, thus altering the functionality of COG-CDG patient neurons.

Smooth transport of cargo by the trafficking machinery is very important during development and synaptic transmission. Defective glycosylation of proteins and lipids disrupts development pathways and alters brain function (Freeze et al., 2012). COG deficient cells also display altered glycosphingolipid biosynthesis. Complex gangliosides are sialic acid containing glycosphingolipids synthesized sequentially, beginning with GM3 and then extended by glycosyltransferases to the more elaborate GM1 gangliosides. Biochemical studies revealed decreased levels of sphingomyelin and GM3 gangliosides in

COG2 deficient CHO cells (Spessott et al., 2010a,b). Gangliosides are ubiquitously expressed, but in the brain the expression of gangliosides and their glycosyltransferases change dramatically during development, from an abundance in of the precursor GM3 to a greater abundance of GM1 (Yu et al., 1988; Kracun et al., 1991, 1992; Ngamukote et al., 2007). Therefore, altered glycosphingolipid biosynthesis could be another reason for the specific neurological manifestation in COG-related congenital disorders.

An additional factor that may add to the apparently higher neuronal vulnerability to COG deficiency is that unlike many other cells, neurons are non-dividing cells and hence cannot easily dilute toxic proteins, peptides, and organelles. Another possibility is that COG is playing a specific and yet uncovered role in neuronal cells. Although the COG complex has been largely detected at the established perinuclear Golgi location in neurons (Figure 1) this does not exclude some sort of moonlighting function for COG subunits at other neuron-specific locations where it could be involved in tethering of specialized vesicles.

Considering the systemic and complex character of COG-related diseases, multiple questions regarding the associated neurological symptoms remain to be addressed. Thus, more *in vivo* studies are needed to dissect the role of COG and its Rab and SNARE partners in these symptoms. Input from COG deficiencies within neuroglia and subtypes of neuroglia presents yet another poorly studied field and avenue requiring further studies. Further research is also needed to establish in which diseases COG-deficiency-associated neurological syndromes are secondary manifestations of some other primary disease state.

## ACKNOWLEDGMENTS

We are thankful to Rachel D. Hendrix and Wezley C. Griffin for critical reading of the manuscript. This work was supported by the NIH grants GM083144 and U54 GM105814, and by the Pilot grant from the Arkansas Biosciences Institute.

## REFERENCES

- Amessou, M., Fradagrada, A., Falguières, T., Lord, J. M., Smith, D. C., Roberts, L. M., et al. (2007). Syntaxin 16 and syntaxin 5 are required for efficient retrograde transport of several exogenous and endogenous cargo proteins. *J. Cell Sci.* 120, 1457–1468. doi: 10.1242/jcs.03436
- Arriagada, C., Bustamante, M., Atwater, I., Rojas, E., Caviedes, R., and Caviedes, P. (2010). Apoptosis is directly related to intracellular amyloid accumulation in a cell line derived from the cerebral cortex of a trisomy 16 mouse, an animal model of Down syndrome. *Neurosci. Lett.* 470, 81–85. doi: 10.1016/j.neulet.2009.12.062
- Bellouze, S., Schäfer, M. K., Buttigieg, D., Baillat, G., Rabouille, C., and Haase, G. (2014). Golgi fragmentation in pmn mice is due to a defective ARF1/TBCE cross-talk that coordinates COPI vesicle formation and tubulin polymerization. *Hum. Mol. Genet.* 23, 5961–5975. doi: 10.1093/hmg/ddu320
- Bonifacino, J. S., and Glick, B. S. (2004). The mechanisms of vesicle budding and fusion. *Cell* 116, 153–166. doi: 10.1016/S0092-8674(03)01079-1
- Burré, J., Sharma, M., Tsetsenis, T., Buchman, V., Etherton, M. R., and Südhof, T. C. (2010). Alpha-synuclein promotes SNARE-complex assembly *in vivo* and *in vitro*. *Science* 329, 1663–1667. doi: 10.1126/science.1195227
- Cataldo, A. M., Peterhoff, C. M., Troncoso, J. C., Gomez-Isla, T., Hyman, B. T., and Nixon, R. A. (2000). Endocytic pathway abnormalities precede amyloid beta deposition in sporadic Alzheimer's disease and down syndrome: differential effects of APOE genotype and presenilin mutations. *Am. J. Pathol.* 157, 277–286. doi: 10.1016/S0002-9440(10)64538-5
- Choudhury, A., Sharma, D. K., Marks, D. L., and Pagano, R. E. (2004). Elevated endosomal cholesterol levels in Niemann-Pick cells inhibit rab4 and perturb membrane recycling. *Mol. Biol. Cell* 15, 4500–4511. doi: 10.1091/mbc.E04-05-0432
- Chua, C. E., and Tang, B. L. (2008). Syntaxin 16 is enriched in neuronal dendrites and may have a role in neurite outgrowth. *Mol. Membr. Biol.* 25, 35–45. doi: 10.1080/09687680701504649
- Cooper, A. A., Gitler, A. D., Cashikar, A., Haynes, C. M., Hill, K. J., Bhullar, B., et al. (2006). Alpha-synuclein blocks ER-Golgi traffic and Rab1 rescues neuron loss in Parkinson's models. *Science* 313, 324–328. doi: 10.1126/science.1129462
- Corbett, M. A., Schwake, M., Bahlo, M., Dibbens, L. M., Lin, M., Gandolfo, L. C., et al. (2011). A mutation in the Golgi Qb-SNARE gene GOSR2 causes progressive myoclonus epilepsy with early ataxia. *Am. J. Hum. Genet.* 88, 657–663. doi: 10.1016/j.ajhg.2011.04.011

- Cottam, N. P., and Ungar, D. (2012). Retrograde vesicle transport in the Golgi. *Protoplasma* 249, 943–955. doi: 10.1007/s00709-011-0361-7
- D'Adamo, P., Masetti, M., Bianchi, V., Morè, L., Mignogna, M. L., Giannandrea, M., et al. (2014). RAB GTPases and RAB-interacting proteins and their role in the control of cognitive functions. *Neurosci. Biobehav. Rev.* 46(Pt 2), 302–314. doi: 10.1016/j.neubiorev.2013.12.009
- De Strooper, B., Iwatsubo, T., and Wolfe, M. S. (2012). Presenilins and gamma-secretase: structure, function, and role in Alzheimer Disease. *Cold Spring Harb. Perspect. Med.* 2:a006304. doi: 10.1101/cshperspect.a006304
- Dugan, J. M., deWit, C., McConlogue, L., and Maltese, W. A. (1995). The Ras-related GTP-binding protein, Rab1B, regulates early steps in exocytic transport and processing of beta-amyloid precursor protein. *J. Biol. Chem.* 270, 10982–10989. doi: 10.1074/jbc.270.18.10982
- Fotso, P., Koryakina, Y., Pavliv, O., Tsiomenko, A. B., and Lupashin, V. V. (2005). Cog1p plays a central role in the organization of the yeast conserved oligomeric Golgi complex. *J. Biol. Chem.* 280, 27613–27623. doi: 10.1074/jbc.M504597200
- Foulquier, F., Ungar, D., Reynders, E., Zeevaert, R., Mills, P., Garcia-Silva, M. T., et al. (2007). A new inborn error of glycosylation due to a Cog8 deficiency reveals a critical role for the Cog1-Cog8 interaction in COG complex formation. *Hum. Mol. Genet.* 16, 717–730. doi: 10.1093/hmg/ddl476
- Foulquier, F., Vasile, E., Schollen, E., Callewaert, N., Raemaekers, T., Quelhas, D., et al. (2006). Conserved oligomeric Golgi complex subunit 1 deficiency reveals a previously uncharacterized congenital disorder of glycosylation type II. *Proc. Natl. Acad. Sci. U.S.A.* 103, 3764–3769. doi: 10.1073/pnas.0507685103
- Freeze, H. H., Chong, J. X., Bamshad, M. J., and Ng, B. G. (2014). Solving glycosylation disorders: fundamental approaches reveal complicated pathways. *Am. J. Hum. Genet.* 94, 161–175. doi: 10.1016/j.ajhg.2013.10.024
- Freeze, H. H., Eklund, E. A., Ng, B. G., and Patterson, M. C. (2012). Neurology of inherited glycosylation disorders. *Lancet Neurol.* 11, 453–466. doi: 10.1016/S1474-4422(12)70040-6
- Fuchs-Telem, D., Stewart, H., Rapaport, D., Nussbeck, J., Gat, A., Gini, M., et al. (2011). CEDNIK syndrome results from loss-of-function mutations in SNAP29. *Br. J. Dermatol.* 164, 610–616. doi: 10.1111/j.1365-2133.2010.10133.x
- Fung, C. W., Matthijs, G., Sturiale, L., Garozzo, D., Wong, K. Y., Wong, R., et al. (2012). COG5-CDG with a mild neurohepatic presentation. *JIMD Rep.* 3, 67–70. doi: 10.1007/8904\_2011\_61
- Ginsberg, S. D., Alldred, M. J., Counts, S. E., Cataldo, A. M., Neve, R. L., Jiang, Y., et al. (2010). Microarray analysis of hippocampal CA1 neurons implicates early endosomal dysfunction during Alzheimer's disease progression. *Biol. Psychiatry* 68, 885–893. doi: 10.1016/j.biopsych.2010.05.030
- Gitler, A. D., Bevis, B. J., Shorter, J., Strathearn, K. E., Hamamichi, S., Su, L. J., et al. (2008). The Parkinson's disease protein alpha-synuclein disrupts cellular Rab homeostasis. *Proc. Natl. Acad. Sci. U.S.A.* 105, 145–150. doi: 10.1073/pnas.0710685105
- Glick, B. S., and Malhotra, V. (1998). The curious status of the Golgi apparatus. *Cell* 95, 883–889. doi: 10.1016/S0092-8674(00)81713-4
- Gonatas, N. K., Stieber, A., and Gonatas, J. O. (2006). Fragmentation of the Golgi apparatus in neurodegenerative diseases and cell death. *J. Neurol. Sci.* 246, 21–30. doi: 10.1016/j.jns.2006.01.019
- Gong, Q., Anderson, C. L., January, C. T., and Zhou, Z. (2002). Role of glycosylation in cell surface expression and stability of HERG potassium channels. *Am. J. Physiol. Heart Circ. Physiol.* 283, H77–H84. doi: 10.1152/ajpheart.00008.2002
- Haas, A. K., Yoshimura, S., Stephens, D. J., Preisinger, C., Fuchs, E., and Barr, F. A. (2007). Analysis of GTPase-activating proteins: Rab1 and Rab43 are key Rabs required to maintain a functional Golgi complex in human cells. *J. Cell Sci.* 120, 2997–3010. doi: 10.1242/jcs.014225
- Hardy, J. (2006). A hundred years of Alzheimer's disease research. *Neuron* 52, 3–13. doi: 10.1016/j.neuron.2006.09.016
- Hasegawa, H., Yang, Z., Olteidal, L., Davanger, S., and Hay, J. C. (2004). Intramolecular protein-protein and protein-lipid interactions control the conformation and subcellular targeting of neuronal Ykt6. *J. Cell Sci.* 117, 4495–4508. doi: 10.1242/jcs.01314
- Hasegawa, H., Zinsner, S., Rhee, Y., Vik-Mo, E. O., Davanger, S., and Hay, J. C. (2003). Mammalian ykt6 is a neuronal SNARE targeted to a specialized compartment by its profilin-like amino terminal domain. *Mol. Biol. Cell* 14, 698–720. doi: 10.1091/mbc.E02-09-0556
- Huybrechts, S., De Laet, C., Bontems, P., Rooze, S., Souayah, H., Sznajder, Y., et al. (2012). Deficiency of subunit 6 of the conserved oligomeric Golgi complex (COG6-CDG): second patient, different phenotype. *JIMD Rep.* 4, 103–108. doi: 10.1007/8904\_2011\_79
- Joshi, G., and Wang, Y. (2015). Golgi defects enhance APP amyloidogenic processing in Alzheimer's disease. *Bioessays* 37, 240–247. doi: 10.1002/bies.201400116
- Kabayama, H., Tokushige, N., Takeuchi, M., and Mikoshiba, K. (2008). Syntaxin 6 regulates nerve growth factor-dependent neurite outgrowth. *Neurosci. Lett.* 436, 340–344. doi: 10.1016/j.neulet.2008.03.061
- Kingsley, D. M., Kozarsky, K. F., Segal, M., and Krieger, M. (1986). Three types of low density lipoprotein receptor-deficient mutant have pleiotropic defects in the synthesis of N-linked, O-linked, and lipid-linked carbohydrate chains. *J. Cell Biol.* 102, 1576–1585. doi: 10.1083/jcb.102.5.1576
- Kodera, H., Ando, N., Yuasa, I., Wada, Y., Tsurusaki, Y., Nakashima, M., et al. (2015). Mutations in COG2 encoding a subunit of the conserved oligomeric golgi complex cause a congenital disorder of glycosylation. *Clin. Genet.* 87, 455–460. doi: 10.1111/cge.12417
- Kracun, I., Rosner, H., Drnovsek, V., Heffer-Laue, M., Cosovic, C., and Laue, G. (1991). Human brain gangliosides in development, aging and disease. *Int. J. Dev. Biol.* 35, 289–295.
- Kracun, I., Rosner, H., Drnovsek, V., Vukelic, Z., Cosovic, C., Trbojevic-Cepe, M., et al. (1992). Gangliosides in the human brain development and aging. *Neurochem. Int.* 20, 421–431. doi: 10.1016/0197-0186(92)90057-X
- Kranz, C., Ng, B. G., Sun, L., Sharma, V., Eklund, E. A., Miura, Y., et al. (2007). COG8 deficiency causes new congenital disorder of glycosylation type IIh. *Hum. Mol. Genet.* 16, 731–741. doi: 10.1093/hmg/ddm028
- Kudlyk, T., Willett, R., Pokrovskaya, I. D., and Lupashin, V. (2013). COG6 interacts with a subset of the Golgi SNAREs and is important for the Golgi complex integrity. *Traffic* 14, 194–204. doi: 10.1111/tra.12020
- Kunwar, A. J., Rickmann, M., Backofen, B., Browski, S. M., Rosenbusch, J., Schöning, S., et al. (2011). Lack of the endosomal SNAREs vti1a and vti1b led to significant impairments in neuronal development. *Proc. Natl. Acad. Sci. U.S.A.* 108, 2575–2580. doi: 10.1073/pnas.1013891108
- Laufman, O., Freeze, H. H., Hong, W., and Lev, S. (2013a). Deficiency of the Cog8 subunit in normal and CDG-derived cells impairs the assembly of the COG and Golgi SNARE complexes. *Traffic* 14, 1065–1077. doi: 10.1111/tra.12093
- Laufman, O., Hong, W., and Lev, S. (2013b). The COG complex interacts with multiple Golgi SNAREs and enhances fusogenic assembly of SNARE complexes. *J. Cell Sci.* 126, 1506–1516. doi: 10.1242/jcs.122101
- Laufman, O., Kedan, A., Hong, W., and Lev, S. (2009). Direct interaction between the COG complex and the SM protein, Sly1, is required for Golgi SNARE pairing. *EMBO J.* 28, 2006–2017. doi: 10.1038/emboj.2009.168
- Lübbelhusen, J., Thiel, C., Rind, N., Ungar, D., Prinsen, B. H., de Koning, T. J., et al. (2010). Fatal outcome due to deficiency of subunit 6 of the conserved oligomeric Golgi complex leading to a new type of congenital disorders of glycosylation. *Hum. Mol. Genet.* 19, 3623–3633. doi: 10.1093/hmg/ddq278
- Lupashin, V., and Ungar, D. (2008). "COG complex," in *The Golgi Apparatus*, eds A. Mironov and M. Pavelka (Vienna: Springer), 120–129. doi: 10.1007/978-3-211-76310-0\_9
- Lupashin, V. V., Pokrovskaya, I. D., McNew, J. A., and Waters, M. G. (1997). Characterization of a novel yeast SNARE protein implicated in Golgi retrograde traffic. *Mol. Biol. Cell* 8, 2659–2676. doi: 10.1091/mbc.8.12.2659
- McConlogue, L., Castellano, F., deWit, C., Schenk, D., and Maltese, W. A. (1996). Differential effects of a Rab6 mutant on secretory versus amyloidogenic processing of Alzheimer's beta-amyloid precursor protein. *J. Biol. Chem.* 271, 1343–1348. doi: 10.1074/jbc.271.3.1343
- Miller, V. J., Sharma, P., Kudlyk, T. A., Frost, L., Roife, A. P., Watson, I. J., et al. (2013). Molecular insights into vesicle tethering at the Golgi by the conserved oligomeric Golgi (COG) complex and the golgin TATA element modulatory factor (TMF). *J. Biol. Chem.* 288, 4229–4240. doi: 10.1074/jbc.M112.426767
- Miller, V. J., and Ungar, D. (2012). Re'COG'nition at the Golgi. *Traffic* 13, 891–897. doi: 10.1111/j.1600-0854.2012.01338.x
- Morava, E., Zeevaert, R., Korsch, E., Huijben, K., Wopereis, S., Matthijs, G., et al. (2007). A common mutation in the COG7 gene with a consistent phenotype including microcephaly, adducted thumbs, growth retardation,



- VSD and episodes of hyperthermia. *Eur. J. Hum. Genet.* 15, 638–645. doi: 10.1038/sj.ejhg.5201813
- Nagahama, M., Orci, L., Ravazzola, M., Amherdt, M., Lacomis, L., Tempst, P., et al. (1996). A v-SNARE implicated in intra-Golgi transport. *J. Cell Biol.* 133, 507–516. doi: 10.1083/jcb.133.3.507
- Nakagomi, S., Barsoum, M. J., Bossy-Wetzel, E., Sütterlin, C., Malhotra, V., and Lipton, S. A. (2008). A Golgi fragmentation pathway in neurodegeneration. *Neurobiol. Dis.* 29, 221–231. doi: 10.1016/j.nbd.2007.08.015
- Ng, B. G., Kranz, C., Hagebeuk, E. E., Duran, M., Abeling, N. G., Wuyts, B., et al. (2007). Molecular and clinical characterization of a Moroccan Cog7 deficient patient. *Mol. Genet. Metab.* 91, 201–204. doi: 10.1016/j.ymgme.2007.02.011
- Ng, B. G., Sharma, V., Sun, L., Loh, E., Hong, W., Tay, S. K., et al. (2011). Identification of the first COG-CDG patient of Indian origin. *Mol. Genet. Metab.* 102, 364–367. doi: 10.1016/j.ymgme.2010.11.161
- Ngamukote, S., Yanagisawa, M., Ariga, T., Ando, S., and Yu, R. K. (2007). Developmental changes of glycosphingolipids and expression of glycogenes in mouse brains. *J. Neurochem.* 103, 2327–2341. doi: 10.1111/j.1471-4159.2007.04910.x
- Paesold-Burda, P., Maag, C., Troxler, H., Foulquier, F., Kleinert, P., Schnabel, S., et al. (2009). Deficiency in COG5 causes a moderate form of congenital disorders of glycosylation. *Hum. Mol. Genet.* 18, 4350–4356. doi: 10.1093/hmg/ddp389
- Pan, P. Y., Cai, Q., Lin, L., Lu, P. H., Duan, S., and Sheng, Z. H. (2005). SNAP-29-mediated modulation of synaptic transmission in cultured hippocampal neurons. *J. Biol. Chem.* 280, 25769–25779. doi: 10.1074/jbc.M502356200
- Parlati, F., Varlamov, O., Paz, K., McNew, J. A., Hurtado, D., Söllner, T. H., et al. (2002). Distinct SNARE complexes mediating membrane fusion in Golgi transport based on combinatorial specificity. *Proc. Natl. Acad. Sci. U.S.A.* 99, 5424–5429. doi: 10.1073/pnas.082100899
- Peric, A., and Annaert, W. (2015). Early etiology of Alzheimer's disease: tipping the balance toward autophagy or endosomal dysfunction? *Acta Neuropathol.* 129, 363–381. doi: 10.1007/s00401-014-1379-7
- Pfeffer, S. R. (2013). Rab GTPase regulation of membrane identity. *Curr. Opin. Cell Biol.* 25, 414–419. doi: 10.1016/j.cob.2013.04.002
- Rendón, W. O., Martínez-Alonso, E., Tomás, M., Martínez-Martínez, N., and Martínez-Menárguez, J. A. (2013). Golgi fragmentation is Rab and SNARE dependent in cellular models of Parkinson's disease. *Histochem. Cell Biol.* 139, 671–684. doi: 10.1007/s00418-012-1059-4
- Reynders, E., Foulquier, F., Annaert, W., and Matthijs, G. (2011). How Golgi glycosylation meets and needs trafficking: the case of the COG complex. *Glycobiology* 21, 853–863. doi: 10.1093/glycob/cwq179
- Reynders, E., Foulquier, F., Leão Teles, E., Quelhas, D., Morelle, W., Rabouille, C., et al. (2009). Golgi function and dysfunction in the first COG4-deficient CDG type II patient. *Hum. Mol. Genet.* 18, 3244–3256. doi: 10.1093/hmg/ddp262
- Richardson, B. C., Smith, R. D., Ungar, D., Nakamura, A., Jeffrey, P. D., Lupashin, V. V., et al. (2009). Structural basis for a human glycosylation disorder caused by mutation of the COG4 gene. *Proc. Natl. Acad. Sci. U.S.A.* 106, 13329–13334. doi: 10.1073/pnas.0901966106
- Rizo, J., and Südhof, T. C. (2012). The membrane fusion enigma: SNAREs, Sec1/Munc18 proteins, and their accomplices—guilty as charged? *Annu. Rev. Cell Dev. Biol.* 28, 279–308. doi: 10.1146/annurev-cellbio-101011-155818
- Rosenbaum, E. E., Vasiljevic, E., Cleland, S. C., Flores, C., and Colley, N. J. (2014). The Gos28 SNARE protein mediates intra-Golgi transport of rhodopsin and is required for photoreceptor survival. *J. Biol. Chem.* 289, 32392–32409. doi: 10.1074/jbc.M114.585166
- Rymen, D., Keldermans, L., Race, V., Régál, L., Deconinck, N., Dionisi-Vici, C., et al. (2012). COG5-CDG: expanding the clinical spectrum. *Orphanet J. Rare Dis.* 7:94. doi: 10.1186/1750-1172-7-94
- Rymen, D., Winter, J., van Hasselt, P. M., Jaeken, J., Kasapkara, C., Gokcay, G., et al. (2015). Key features and clinical variability of COG6-CDG. *Mol. Genet. Metab.* doi: 10.1016/j.ymgme.2015.07.003. [Epub ahead of print].
- Saito, T., Suemoto, T., Brouwers, N., Slegers, K., Funamoto, S., Mihira, N., et al. (2011). Potent amyloidogenicity and pathogenicity of Abeta43. *Nat. Neurosci.* 14, 1023–1032. doi: 10.1038/nn.2858
- Schardt, A., Brinkmann, B. G., Mitkovski, M., Sereda, M. W., Werner, H. B., and Nave, K. A. (2009). The SNARE protein SNAP-29 interacts with the GTPase Rab3A: implications for membrane trafficking in myelinating glia. *J. Neurosci. Res.* 87, 3465–3479. doi: 10.1002/jnr.22005
- Scheper, W., Zwart, R., and Baas, F. (2004). Rab6 membrane association is dependent of Presenilin 1 and cellular phosphorylation events. *Brain Res. Mol. Brain Res.* 122, 17–23. doi: 10.1016/j.molbrainres.2003.11.013
- Scott, H., and Panin, V. M. (2014). N-glycosylation in regulation of the nervous system. *Adv. Neurobiol.* 9, 367–394. doi: 10.1007/978-1-4939-1154-7\_17
- Shaheen, R., Ansari, S., Alshammari, M. J., Alkhalidi, H., Alrukban, H., Eyaad, W., et al. (2013). A novel syndrome of hypohidrosis and intellectual disability is linked to COG6 deficiency. *J. Med. Genet.* 50, 431–436. doi: 10.1136/jmedgenet-2013-101527
- Shestakova, A., Suvorova, E., Pavliv, O., Khaidakova, G., and Lupashin, V. (2007). Interaction of the conserved oligomeric Golgi complex with t-SNARE Syntaxin5a/Sed5 enhances intra-Golgi SNARE complex stability. *J. Cell Biol.* 179, 1179–1192. doi: 10.1083/jcb.200705145
- Shestakova, A., Zolov, S., and Lupashin, V. (2006). COG complex-mediated recycling of Golgi glycosyltransferases is essential for normal protein glycosylation. *Traffic* 7, 191–204. doi: 10.1111/j.1600-0854.2005.00376.x
- Siddiqi, S., Mani, A. M., and Siddiqi, S. A. (2010). The identification of the SNARE complex required for the fusion of VLDL-transport vesicle with hepatic cis-Golgi. *Biochem. J.* 429, 391–401. doi: 10.1042/BJ20100336
- Smith, R. D., Willett, R., Kudlyk, T., Pokrovskaya, I., Paton, A. W., Paton, J. C., et al. (2009). The COG complex, Rab6 and COPI define a novel Golgi retrograde trafficking pathway that is exploited by SubAB toxin. *Traffic* 10, 1502–1517. doi: 10.1111/j.1600-0854.2009.00965.x
- Spaapen, L. J. M., Bakker, J. A., van der Meer, S. B., Sijstermans, H. J., Steet, R. A., Wevers, R. A., et al. (2005). Clinical and biochemical presentation of siblings with COG-7 deficiency, a lethal multiple O- and N-glycosylation disorder. *J. Inher. Metab. Dis.* 28, 707–714. doi: 10.1007/s10545-005-0015-z
- Spessott, W., Uliana, A., and Maccioni, H. J. (2010a). Cog2 null mutant CHO cells show defective sphingomyelin synthesis. *J. Biol. Chem.* 285, 41472–41482. doi: 10.1074/jbc.M110.150011
- Spessott, W., Uliana, A., and MacCioni, H. J. (2010b). Defective GM3 synthesis in Cog2 null mutant CHO cells associates to mislocalization of lactosylceramide sialyltransferase in the Golgi complex. *Neurochem. Res.* 35, 2161–2167. doi: 10.1007/s11064-010-0319-8
- Sprecher, E., Ishida-Yamamoto, A., Mizrahi-Koren, M., Rapaport, D., Goldsher, D., Indelman, M., et al. (2005). A mutation in SNAP29, coding for a SNARE protein involved in intracellular trafficking, causes a novel neurocutaneous syndrome characterized by cerebral dysgenesis, neuropathy, ichthyosis, and palmoplantar keratoderma. *Am. J. Hum. Genet.* 77, 242–251. doi: 10.1086/432556
- Steggmaier, M., Yang, B., Yoo, J. S., Huang, B., Shen, M., Yu, S., et al. (1998). Three novel proteins of the syntaxin/SNAP-25 family. *J. Biol. Chem.* 273, 34171–34179. doi: 10.1074/jbc.273.51.34171
- Steet, R., and Kornfeld, S. (2006). COG-7-deficient human fibroblasts exhibit altered recycling of golgi proteins. *Mol. Biol. Cell* 17, 2312–2321. doi: 10.1091/mbc.E05-08-0822
- Subramaniam, V. N., Peter, F., Philp, R., Wong, S. H., and Hong, W. (1996). GS28, a 28-kilodalton Golgi SNARE that participates in ER-Golgi transport. *Science* 272, 1161–1163. doi: 10.1126/science.272.5265.1161
- Suga, K., Hattori, H., Saito, A., and Akagawa, K. (2005a). RNA interference-mediated silencing of the syntaxin 5 gene induces Golgi fragmentation but capable of transporting vesicles. *FEBS Lett.* 579, 4226–4234. doi: 10.1016/j.febslet.2005.06.053
- Suga, K., Saito, A., and Akagawa, K. (2015). ER stress response in NG108-15 cells involves upregulation of syntaxin 5 expression and reduced amyloid beta peptide secretion. *Exp. Cell Res.* 332, 11–23. doi: 10.1016/j.yexcr.2015.01.001
- Suga, K., Saito, A., Tomiyama, T., Mori, H., and Akagawa, K. (2005b). Syntaxin 5 interacts specifically with presenilin holoproteins and affects processing of betaAPP in neuronal cells. *J. Neurochem.* 94, 425–439. doi: 10.1111/j.1471-4159.2005.03210.x
- Thayanidhi, N., Helm, J. R., Nycz, D. C., Bentley, M., Liang, Y., and Hay, J. C. (2010). Alpha-synuclein delays endoplasmic reticulum (ER)-to-Golgi transport in mammalian cells by antagonizing ER/Golgi SNAREs. *Mol. Biol. Cell* 21, 1850–1863. doi: 10.1091/mbc.E09-09-0801
- Ungar, D., Oka, T., Vasile, E., Krieger, M., and Hughson, F. M. (2005). Subunit architecture of the conserved oligomeric Golgi complex. *J. Biol. Chem.* 280, 32729–32735. doi: 10.1074/jbc.M504590200



- von Mollard, G. F., Nothwehr, S. F., and Stevens, T. H. (1997). The yeast v-SNARE Vti1p mediates two vesicle transport pathways through interactions with the t-SNAREs Sed5p and Pep12p. *J. Cell Biol.* 137, 1511–1524. doi: 10.1083/jcb.137.7.1511
- Walter, A. M., Kurps, J., de Wit, H., Schoning, S., Toft-Bertelsen, T. L., Lauks, J., et al. (2014). The SNARE protein vti1a functions in dense-core vesicle biogenesis. *EMBO J.* 33, 1681–1697. doi: 10.15252/embj.201387549
- Watanabe, I., Zhu, J., Recio-Pinto, E., and Thornhill, W. B. (2004). Glycosylation affects the protein stability and cell surface expression of Kv1.4 but not Kv1.1 potassium channels. A pore region determinant dictates the effect of glycosylation on trafficking. *J. Biol. Chem.* 279, 8879–8885. doi: 10.1074/jbc.M309802200
- Whyte, J. R., and Munro, S. (2002). Vesicle tethering complexes in membrane traffic. *J. Cell Sci.* 115, 2627–2637.
- Willett, R., Kudlyk, T., Pokrovskaya, I., Schönherr, R., Ungar, D., Duden, R., et al. (2013a). COG complexes form spatial landmarks for distinct SNARE complexes. *Nat. Commun.* 4, 1553. doi: 10.1038/ncomms2535
- Willett, R., Pokrovskaya, I., Kudlyk, T., and Lupashin, V. (2014). Multipronged interaction of the COG complex with intracellular membranes. *Cell. Logist.* 4:e27888. doi: 10.4161/cl.27888
- Willett, R., Ungar, D., and Lupashin, V. (2013b). The Golgi puppet master: COG complex at center stage of membrane trafficking interactions. *Histochem. Cell Biol.* 140, 271–283. doi: 10.1007/s00418-013-1117-6
- Wu, X., Steet, R. A., Bohorov, O., Bakker, J., Newell, J., Krieger, M., et al. (2004). Mutation of the COG complex subunit gene COG7 causes a lethal congenital disorder. *Nat. Med.* 10, 518–523. doi: 10.1038/nm1041
- Xu, Y., Martin, S., James, D. E., and Hong, W. (2002). GS15 forms a SNARE complex with syntaxin 5, GS28, and Ykt6 and is implicated in traffic in the early cisternae of the Golgi apparatus. *Mol. Biol. Cell* 13, 3493–3507. doi: 10.1091/mbc.E02-01-0004
- Yu, R. K., Macala, L. J., Taki, T., Weinfield, H. M., and Yu, F. S. (1988). Developmental changes in ganglioside composition and synthesis in embryonic rat brain. *J. Neurochem.* 50, 1825–1829. doi: 10.1111/j.1471-4159.1988.tb02484.x
- Zeevaert, R., Foulquier, F., Cheillan, D., Cloix, I., Guffon, N., Sturiale, L., et al. (2009a). A new mutation in COG7 extends the spectrum of COG subunit deficiencies. *Eur. J. Med. Genet.* 52, 303–305. doi: 10.1016/j.ejmg.2009.06.006
- Zeevaert, R., Foulquier, F., Dimitrov, B., Reyniers, E., Van Damme-Lombaerts, R., Simeonov, E., et al. (2009b). Cerebrocostomandibular-like syndrome and a mutation in the conserved oligomeric Golgi complex, subunit 1. *Hum. Mol. Genet.* 18, 517–524. doi: 10.1093/hmg/ddn379
- Zhang, T., and Hong, W. (2001). Ykt6 forms a SNARE complex with syntaxin 5, GS28, and Bet1 and participates in a late stage in endoplasmic reticulum-Golgi transport. *J. Biol. Chem.* 276, 27480–27487. doi: 10.1074/jbc.M102786200
- Zolov, S. N., and Lupashin, V. V. (2005). Cog3p depletion blocks vesicle-mediated Golgi retrograde trafficking in HeLa cells. *J. Cell Biol.* 168, 747–759. doi: 10.1083/jcb.200412003

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Climer, Dobretsov and Lupashin. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.